



Original Research Article

An investigation into the Anti-inflammatory Activity of n-hexane Extract of the Leaves of *Baphia pubescens* Hook. F (Family Leguminosae)

Anowi Chinedu Fredrick*¹, Uzodinma S², Utoh-Nedosa UA³, Onyegbule Felix A⁴

¹ Department of Pharmacognosy and Traditional Medicine, Faculty of Pharm Sciences, Nnamdi Azikiwe University, Awka, Nigeria

² Department of Clinical Pharmacy and Pharmacy Management, Faculty of Pharm Sciences, Nnamdi Azikiwe University, Awka, Nigeria

³ Department of Pharmacology and Toxicology, Faculty of Pharm Sciences, Nnamdi Azikiwe University, Awka, Nigeria

⁴ Department of Medicinal and Pharmaceutical Chemistry, Faculty of Pharm Sciences, Nnamdi Azikiwe University, Awka, Nigeria

* Corresponding Author

Received: 03 July 2014

Revised: 11 July 2014

Accepted: 21 July 2014

ABSTRACT

Baphia pubescens is claimed by the people of Ogidi in Idemili Local Government area of Anambra State, Nigeria to have anti pyretic, anti-inflammatory, analgesic and antimicrobial properties. The decoction of the leaves is used to treat wounds, running stomach (diarrhea), aches and pains as well as fever locally. This investigation was carried out to ascertain the veracity of these claims.

The leaves of *Baphia pubescens* was collected and dried at ambient temperature and pulverized. Exactly 200 g of the powdered drug was extracted with 400ml of n-hexane using the cold maceration technique for 24hours with occasional shaking. This was filtered and the procedure repeated with the marc. The combined filtrates were concentrated under reduced pressure with rotary evaporator. The preliminary phytochemical tests were carried out using standard methods. The anti-inflammatory activity was induced by egg albumin and anti-inflammation measured at predetermined time after the administration of the drug.

The leaves of *Baphia pubescens* exhibited anti-inflammatory activity by reducing the inflammation in rats. Alkaloids, flavonoids, saponins carbohydrates, carbohydrates were absent while steroids and tannins, were found. Preliminary studies support the claim that the leaves of *Baphia pubennscens* possess anti-inflammatory properties hence it is used by the native as anti-inflammatory agent.

Keywords: *Baphia pubescens*; marc; brewer's yeast; rotary evaporator

INTRODUCTION

Herbal medicine practice plays an important role in the primary healthcare delivery system in most

developing countries including Nigeria [1]. Even the World Health Organization (WHO, 2002) is actively encouraging national governments of member

countries to utilize their traditional systems of medicines with regulations suitable to their national health care systems. The WHO estimates that 80% of the population living in rural areas use or depend on herbal medicine for their health needs (WHO Traditional Medicine Strategy, 2002). However, in spite of the obvious and important contribution the herbal medicine makes to primary health care, it continues to be antagonised by majority of allopathic medical practitioners as it is considered to have no scientific basis [2]. This work is therefore a preliminary work to prove that there is scientific evidence to the use of leaves of *Baphia pubescens* in the treatment of pyrexia.

One major problem of herbal medicine practice is that there is no official standard and / or local monograph [3]. In Nigeria, the Federal Government has urged the federating states to set up traditional medicine boards to license and regulate the practice of herbal practitioners under the supervision of ministries of health [4].

Many medicines including reserpine, ergotamine, vincristine, and vinblastine are of herbal origin [5]. About one quarter of the present prescription drugs dispensed by community pharmacies in the United States contain at least one active principle originally derived from plant materials.

***Baphia Pubescens*: Plant Profile**

Taxon: *Baphia pubescens* Hook.F.

Genus: *Baphia*

Family: Leguminosae -papillonoideae

Tribe : Sophoreae

Synonym: *Baphia bancoensis* Aubrev

Common names:

English names: Benin camwood

Other names: Awewi, urohun, Maajigi

Description of the Plants

A tree about 20 ft. high, with trunk 20cm in diameter the ultimate branches slender, terete, densely brown-silky. Petioles 3/8– 1/2 in. long,

slender, ferruginous; leaves oblong or narrow-obovate, 3–4 in. long, acuminate, the base cuneate or slightly rounded, subcoriaceous, under surface ferruginous on the veins when young. Pedicels 1–4 together from the main branches, 1/4 in. long, erecto-patent, ferruginous-downy. Bracteoles minute, rounded. Calyx 1/4 in. deep, finely ferruginous-downy. Corolla twice as long as the calyx, white; standard roundish, 1/2 in. broad. Pod straight, 3 in. long, 3/4 in. broad, membranous, rigid, glabrous, brown, polished, narrowed to both ends [6].



Geographical Distribution

Benin camwood (*Baphia pubescens* Hook.F) - It has a distribution similar to that of *Baphia nitida*, It's main geographical area is Africa found in countries including Nigeria, Zarie, Congo, Ivory Coast , Benin, Cameroon, Gabon, Ghana, Liberia [7].

Habitat: Guineo- Congolian forest; Guinea Congolia/Sudania regional transition zone forest [8]

Medicinal Uses

The leaves or leaf juice are applied against parasitic skin diseases. A leaf infusion is drunk to cure enteritis and other gastrointestinal problems [9].

In Ghana, Côte d'Ivoire and Nigeria the leaves and bark are considered haemostatic and anti-inflammatory, and are used for healing sores and wounds [10].

In Côte d'Ivoire powdered leaves are taken with palm wine or food to cure venereal diseases, and leaf sap is applied as eye drops against jaundice [11].

An extract of young leaves with some salt and red pepper is used as nose drops against headache [12].

In Nigeria powdered heartwood is made into an ointment with shea butter (obtained from the seeds of *Vitellaria paradoxa* C.F.Gaertn.) which is applied against stiff and swollen joints, sprains and rheumatic complaints [13].

In Sierra Leone a bark decoction is drunk to cure cardial pain and bark and leaves are prepared as an enema to treat constipation [14].

In Nigeria and Ghana the pounded dried root, mixed with water and oil, is applied to a ringworm-like fungus attack

In Côte d'Ivoire a leaf extract of camwood and *Senna occidentalis* (L.) Link is drunk against asthma.

In Benin a decoction of the leaves is taken against jaundice and diabetes; in combination with leaves of *Morinda lucida* Benth. it is a treatment against female sterility and painful menstruation. In the leaves saponins, flavonoids, glycosides and

true tannins are present. An ointment made from the leaves showed anti-inflammatory activity in mice. Extracts of fresh leaves inhibited digestion in mice and rats, and showed anti-diarrhea

Leaf extracts of *Baphia nitida* have also been found to show analgesic effect in mice [13].

MATERIALS AND METHOD

Materials

N-hexane, tween-80, Paracetamol , ibuprofen and Aspirin tablets.

Collection and Identification

The fresh leaves of *Baphia pubescens* were obtained from Ogidi, Idemili North Local Government Area of Anambra State, Nigeria in December 2013, during the dry season and was identified by Mr Ozioko, a Taxonomist with the Biosource Development and Conservation program(BDCP) Nsukka, Enugu State, Nigeria. The leaves were air-dried for 2 weeks in the Pharmacognosy laboratory. They were milled and 500g of the powdered plant material was obtained.

Preparation of Ethanol Extract

200g of powered plant sample was macerated in 400mls of n-hexane, for 48 hours after which it was filtered with muslim cloth and further filtered using Whatman (No 1) filter paper. The procedure was repeated with the marc. The combined filtrates were concentrated using rotary 45°C.

Phytochemical Screening

The tests carried out ware based on procedures outlined by [15] and [16].

Tests for Alkaloids

To 0.5gm of the extract, 5.0 ml of 1% aqueous hydrochloric acid was added steamed on a steam bath and filtered. 1.0 ml of the filtrate was then treated with five drops of Mayer's reagent and a second 1.0 ml portion treated similarly with freshly prepared Dragendorffs and Wagner's reagents.

Turbidity or precipitations with either of the reagents indicated the presence of alkaloids in the extract.

Test for Tannins

To 0.5g of the extract, 20ml of water was added, boiled and, filtered and used for the following test

(a) Ferric chloride test

To 3ml of the filtrate, 2 drops of ferric chloride was added. Formation of a greenish black precipitate indicated the presence of tannins.

(b) Lead acetate test

To 3ml of the filtrate was added lead acetate solution. Formation of precipitate indicated the presence of tannins

Test for Saponins

To 1.0gm of the plant extract, 5.0 ml of distilled water was added. The solution was shaken in a test tube and filtered. Frothing which persists on warming is a preliminary evidence for the presence of saponins. 10.0 ml of the filtrate was mixed with 5.0 ml of distilled water and shaken vigorously for a stable persistent froth. The frothing was mixed with 3 drops of olive oil and shaken vigorously then observed for the formation of emulsion which confirms the presence of saponins

Test for Proteins

To 0.5g of the extract, 20ml of distilled water was added, shaken and, filtered and the filtrate was used for the following tests

(a) Millon's test

To a little portion of the filtrate in a test tube, two drops of millon's reagent was added. A white precipitate indicated the presence of proteins.

(b) Xanthoproteic test

5ml of the filtrate was heated with few drops of concentrated nitric acid. A yellow colour which changed to orange on addition of an alkali (dilute sodium hydroxide) indicated the presence of protein.

Test for Flavonoids

10ml of ethyl acetate was added to 0.2g of the extract and heated on a water bath for 3 minutes. The mixture was cooled, filtered and the filtrate was used for the following tests

(a) Ammonium test

4ml of the filtrate was shaken with 1ml of dilute ammonia solution. The layers were allowed to separate. A yellow colour in the ammoniacal layer indicated the presence of flavonoids.

(b) 1% Aluminium chloride solution test

4ml of the filtrate was shaken with 1ml of 1% aluminium chloride solution and the layers were allowed to separate. The formation of yellow colour in the aluminium chloride layer indicated the presence of flavonoids

Test for Steroids and Terpenoids

Exactly 9ml of ethanol was added to 1g of the extract and refluxed for a few minutes and filtered. The filtrate was concentrated to 2.5ml on a boiling water bath. 5ml of hot water was added to the concentrated solution, the mixture was allowed to stand for 1 hour and the waxy matter was filtered off. The filtrate was extracted with 2.5ml of chloroform using separating funnel.

To 0.5ml of the chloroform extract in a test tube was carefully added 1ml of concentrated sulphuric acid to form a lower layer. A reddish brown interface showed the presence of steroids.

Another 0.5ml of the chloroform extract was evaporated to dryness on a water bath and heated with 3ml of concentrated sulphuric acid for 10 minutes on a water bath. A grey colour indicates the presence of terpenoids.

Test for Carbohydrates

Molisch test

To 0.1g of the extract 2ml of water was added, boiled, and filtered. To the filtrate, two drops of naphthol solution in ethanol (molisch reagent) was

added. Concentrated sulphuric acid was gently poured down the side of the test tube to form a lower layer. A purple interfacial ring indicated the presence of carbohydrate.

Test for reducing sugars

0.1g of the plant extract was shaken vigorously with 5ml of distilled water and filtered. The filtrate was divided and used as follows

Fehling's test

To a 1ml portion of the filtrate was an added equal volume of fehing's solution I and II and boiled on a water bath for a few minutes. A brick red precipitate indicated the presence of reducing sugars.

Animals

White male albino rats (150-250kg) obtained from the animal house of the Department of Pharmacology and Toxicology of Madonna university, Elele Campus, River state were used for this study. All the animals were housed under standard environmental conditions where they have free access to food and water.

Anti- Inflammatory Activity

The egg albumin method [17] was employed for this test. 15 rats were used for this study ie 3 rats in each group for four groups. The rats in the first 3 groups were given 100mg/kg, 200mg/kg and 400mg/kg of the extract orally respectively. Group 4 rats were given 1ml of tween 80 as negative control and group 5 rats were given 100mg of ibuprofen as standard drug. After 45mins of administration, inflammation was induced by subcutaneous administration of 0.1ml egg albumin in the right leg of the rats; the rate of swelling was measured in cm at 1hr, 2hrs, 3hrs and 4hours.

Statistical Analysis

All procedures were carried out in triplicates and the results expressed as \bar{x} ±standard error of mean (SEM).

RESULT AND DISCUSSION

Results of phytochemical tests are represented in table 1. After 4hrs of the administration of the drugs, the dose of 400mg/kg exhibited activity almost equivalent to the standard drug. At 4hrs, the activity of the dose of 200mg/kg was almost equal to the standard drug. The drugs appeared to have fast onset of action. Hence after 1hr of the administration of the drug, the 200mg/kg was equivalent to the standard drug whereas the 400mg/kg at this time was more effective (table 2).

Table 1: Phytochemical Result

Constituents	Relative Appearance
Saponins	-
Tannins	+
Carbohydrates	-
Protein	-
Flavonoids	-
Reducing sugar	-
Alkaloids	-
Steroids	++

Table 2: Anti-Inflammatory Result

	0hr	1hr	2hrs	3hrs	4hrs
100mg	2.0±	4.1±	4.1±	3.9±0.	3.8±
/kg	0.00	0.04	0.04	03	0.03
200mg	2.0±	3.9±	3.8±	2.9±	2.9
/kg	0.00	0.02	0.03	0.02	±0.03
400mg	2.0±	3.8±	3.6±	2.5±	2.4±
/kg	0.00	0.02	0.00	0.02	0.02
Control	1.9±0.	4.1±0.	4.3±0.	4.0±0.	3.9±0.
	04	02	02	02	02
Standa	2.0±0.	3.9±0.	3.5±0.	2.6±0.	2.2±0.
rd	00	03	02	00	02

Many reviews and articles reporting the biological activities of flavonoids [18] anthraquinones, polyphenols and phenols, and tannins [19] have been published in recent years. Several phenol compounds have been identified and isolated from plants and they have shown promising bacterial inhibiting properties against specific and broad

spectrum of cultured as well as clinical bacterial strains including Methicillin-Resistant *Staphylococcus aureus* (MRSA), and multi-drug resistant bacteria. The presence of alkaloids has been shown to demonstrate biological activity [20]. Alkaloids, phenols, flavonoids and glycosides have a number of biological activities and strong antibacterial potentials. Alkaloids have exhibited promising activity against *H. pylori* [21] and a number of other bacterial strains [22-23]. The Result of phytochemical screening showed abundance of steroids and moderate amount of tannins but absence of carbohydrates, alkaloids, saponins, flavonoids, proteins and reducing sugars in the n-hexane extract of *Baphia pubescens*.

CONCLUSION

The n-hexane leaves extract of *Baphia pubescens* exhibited anti-inflammatory activity; hence it is used by the local community in Ogidi of Anambra State, Nigeria as anti-inflammatory drug.

REFEERNCES

- Hostettmann K. Marston, a Study in Natural Products Chemistry. Amsterdam: Elsevier; 1990, p 405.
- Odugbemi T. A Textbook of Medicinal Plants from Nigeria. Lagos: University of Lagos Press; 2008.
- Nadkarn K M. Indian plants with their medicinal properties and uses. Indian Journal of Chemistry 2001; 48: 282–293.
- Allen ASK, Reisch J. Phytochemistry, African Pharmacopoeia, Vol. 1, STRC/OAU: Lagos, 1985, 24, 3003.
- Ajaiyeoba E O. Phytochemical and antimicrobial studies of *Gynandropsis gynandra* and *Buchholzia Coriaceae* extracts. African Journal of Biomedical Research 2000; 3: 161 – 165.
- Craig W J, (1999) Health-promoting properties of common herbs. American Journal of Clinical Nutrition 1999; 70: 3: 491–499.
- Timothy SY, Wazis CH, Zakama SG, Dawurung JS and Albert T. Antipyretic activity of aqueous and ethanolic extract of *Casia alata* Linn Leaf. *International Journal of Research in Ayurveda & Pharmacy* 2012; 3(6): 811-813.
- Sofowora A. Medicinal plants and traditional medicine in Africa. Ibadan, Nigeria: Spectrum Books Ltd; 1993, p 289.
- FAO. Some medicinal forest plants of Africa and Latin America. *Journal of Science* 1996; 67: 75–78,161–162.
- Akande IS, Akande BE and Gbenle GO. Toxicology and antioxidant effects of *Baphia nitida* on diazepam Induced oxidative stress in rats. *African Journal of Biochemistry Research* 2011; 5(8):255-263.
- Konkon ND, Quattara D, Adjoungoua AL, Simaga AL and Kone B. Anti-inflammatory and cicatrizing activity of *Baphia nitida* exudates. *International Journal of Biosciences (IJB)* 2011; 1(5):45-50.
- Onwukaeme ND. Anti- inflammatory activities of flavonoids of *Baphia nitida* Lodd. (Leguminosae) on mice and rats. *Journal of Ethnopharmacology* 1995; 46(2): 121-124.
- Mbranchta M, Kamanyi A, Teponno R, Tapondjou A, Watcho P and Nguelefack T. Analgesic and anti-inflammatory properties of extracts from the bulbils of *Dioscorea bulbifera* L. Var sativa (dioscoreaceae) in mice and rats. *Evidence-Based Complementary and Alternative Medicine (eCAM)* 2011; 1-9.
- Kumar KT, Gorain B, Roy DK, Zothanpuia Z, Samanta SK, Pal M, Biswas P, Roy A, Adhikari D, Karmakar S and Sen T. Anti-inflammatory activity of *Acanthus ilicifolius*. *Journal of Ethnopharmacology* 2008; 120: 7-12.
- Harbourne JB. Phytochemical methods: A guide to modern technique of plant analysis. London: Chapman and Hall; 1984.
- Trease GE and Evans WC. Pharmacognosy. London: Saunders Publishers; 2002.

17. Krishna Kumari T, Packia Lincy M, Muthukumarasamy S and Mohan VR. Anti-inflammatory activity of *Sacostemma secamone* (L.) Bennet. *Bioscience Discovery* 2012; 3(3): 288-291.
18. Pretorius JC. Antiinfective agents. *Current Medicinal Chemistry* 2003; 2: 335-353.
19. Tominaga K et al. In vivo action of novel alkyl methyl quinolone alkaloids against *Helicobacter pylori*. *Journal of Antimicrobial Chemotherapy* 2002; 50:547-552.
20. Gianina et al. The antimicrobial effects of malunggay root extract on *E. coli*, *S. aureus* and *Candida albicans*. 2003. Accessed at: <http://www.scinet.dost.gov.ph> on November 6, 2012.
21. Hadi S and Bremner JB. Chemistry, biological and pharmacological properties of African medicinal plants. Harare: University of Zimbabwe Publications; 1996.
22. Khan MR, Kihara M and Omoloso AD. Antimicrobial activity of *Clematis papuasica* and *Nauclea obversifolia*. *Fitoterapia* 2001; 72: 575-578.
23. Kren V and Martinkova L. Glycosides in medicine: the role of glycosidic residue in biological activity. *Current Medicinal Chemistry* 2001; 8:1303-1328.

Cite this article as:

Anowi Chinedu Fredrick, Uzodinma S , Utoh-Nedosa UA and Onyegbule Felix A. An investigation into the anti-inflammatory activity of n-hexane extract of the leaves of *Baphia pubescens hook. F* (family leguminosae). *J Pharm Chem Biol Sci* 2014; 2(2):104-110.